Please check the examination details below	ow before enter	ring your candid	ate information
Candidate surname		Other names	OS:/British
Centre Number Candidate Nu	umber		John de Harman d
Pearson Edexcel Inter	nationa	al Adva	nced Level
<b>Time</b> 1 hour 30 minutes	Paper reference	WCH	H11/01
Chemistry			
International Advanced Su	ubsidiary	//Advanc	ed Level
UNIT 1: Structure, Bondir Organic Chemistry	ng and In	troduction	on to
You must have: Scientific calculator, ruler			Total Marks

# **Instructions**

- Use **black** ink or **black** ball-point pen.
- Fill in the boxes at the top of this page with your name, centre number and candidate number.
- Answer all questions.
- Answer the questions in the spaces provided
  - there may be more space than you need.

### Information

- The total mark for this paper is 80.
- The marks for **each** question are shown in brackets
  - use this as a guide as to how much time to spend on each question.
- You will be assessed on your ability to organise and present information, ideas, descriptions and arguments clearly and logically, including your use of grammar, punctuation and spelling.
- A Periodic Table is printed on the back cover of this paper.

## **Advice**

- Read each question carefully before you start to answer it.
- Show all your working in calculations and include units where appropriate.
- Check your answers if you have time at the end.

Turn over ▶







SECTION A

Answer ALL the questions in this section.

You should aim to spend no more than 20 minutes on this section.

For each question, select one answer from A to D and put a cross in the box . If you change your mind, put a line through the box  $\boxtimes$  and then mark your new answer with a cross  $\boxtimes$ .

- The maximum permitted concentration of sulfur in diesel fuel is 10 mg of sulfur in 1 kg of diesel fuel.
  - (a) What is this concentration of sulfur in ppm?

(1)

- **A** 0.00001
- X 0.01 В
- C 10
- X **D** 10000
- (b) 3.2 kg of this diesel fuel is burned in air.

What is the maximum volume, in dm<sup>3</sup>, of sulfur dioxide which can be produced, measured at room temperature and pressure (r.t.p.)?

[Molar volume of a gas at r.t.p. =  $24 \,\mathrm{dm}^3 \,\mathrm{mol}^{-1}$ ]

(1)


- 0.024
- X В 0.77
- X C 2.4
- X **D** 24

(Total for Question 1 = 2 marks)

- Which equation represents the **second** ionisation energy of chlorine?
  - X **A**  $Cl^{-}(q) + e^{-}$  $\rightarrow$  Cl<sup>2-</sup>(a)
  - $+ 2e^{-} \rightarrow Cl^{2-}(q)$ X **B** Cl(q)
  - $2Cl(g) \rightarrow 2Cl^{+}(g) +$ 2e<sup>-</sup>
  - $Cl^+(g) \rightarrow Cl^{2+}(g) + e^-$ X D

(Total for Question 2 = 1 mark)

The diagram shows a mass spectrometer.



(a) Which process occurs in region **R**?

**A** the sample is vaporised using a heater

- X electrons are removed from molecules or atoms and positive ions are formed
- X **C** electrons are added to the molecules or atoms and negative ions are formed
- X **D** ions are accelerated by an electric field
- (b) Which statement is correct for region **T**?

(1)

(1)

- X **A** ions with a greater mass have a smaller deflection
- X ions with a greater mass have a greater deflection
- X **C** ions with a greater charge have a smaller deflection
- X **D** ions are speeded up by a magnetic field

(Total for Question 3 = 2 marks)

A mass of 0.23 g of sodium was added to  $350\,\mathrm{cm^3}$  water to form hydrogen and a solution of sodium hydroxide.  $^{112(c)} + H_2O(l) \rightarrow NaOH(aq) + \frac{1}{2}H_2(g)$  Independent of the solution of the solution of sodium hydroxide in the

Na(s) + 
$$H_2O(l) \rightarrow NaOH(aq) + \frac{1}{2}H_2(g)$$

- X **A** 0.010
- X **B** 0.029
- X **C** 0.29
- X **D** 0.66
- (b) What is the maximum volume, in cm<sup>3</sup>, of hydrogen which could be formed, measured at r.t.p.?

[Molar volume of a gas at r.t.p. =  $24 \,\mathrm{dm^3} \,\mathrm{mol^{-1}}$ ]

(1)

- X 120
- X 240 В
- X C 480
- X **D** 2800
- (c) The sodium hydroxide solution was neutralised with sulfuric acid.

Which is the ionic equation for this reaction?

(1)

- X **A**  $H^+(aq) + OH^-(aq) \rightarrow H_2O(l)$
- $\times$ **B**  $SO_4^{2-}(aq) + 2Na^+(aq) \rightarrow Na_2SO_4(aq)$
- X **C**  $H_2SO_4(aq) + 2Na^+(aq) + 2OH^-(aq) \rightarrow Na_2SO_4(aq) + 2H_2O(l)$
- X **D**  $2H^{+}(aq) + SO_{4}^{2-}(aq) + 2Na^{+}(aq) + 2OH^{-}(aq) \rightarrow 2Na^{+}(aq) + SO_{4}^{2-}(aq) + 2H_{2}O(l)$

(d) Sodium hydroxide solution was added to magnesium sulfate solution.

The equation for the reaction is shown.

$$2NaOH(aq) \ + \ MgSO_4(aq) \ \rightarrow \ Mg(OH)_2(s) \ + \ Na_2SO_4(aq)$$

hiths:/hritishshidentroom.bd30a.web.app. What is the atom economy (by mass) for the production of magnesium hydroxide?

[A<sub>r</sub> values: H = 1.0O = 16.0Na = 23.0Mg = 24.3S = 32.1

(1)

- X 29.1%
- X 41.0%
- X 48.4%
- X 50.0%

(Total for Question 4 = 4 marks)



- 5 In which series are the ions in order of **decreasing** ionic radius?
  - $\square$  **A**  $Al^{3+}$  >  $Mg^{2+}$  >  $Na^{+}$
  - $\square$  **B** Li<sup>+</sup> > Na<sup>+</sup> > K<sup>+</sup>
  - $\square$  **C**  $N^{3-} > O^{2-} > F^{-}$
  - $\square$  **D**  $O^{2-}$  >  $S^{2-}$  >  $Se^{2-}$

radius?

Total for Question 5 = 1 mark)

**6** A stable ion, **M**<sup>3+</sup>, contains 18 electrons.

In which block of the Periodic Table is element **M** found?

- $\mathbf{X}$  A  $\mathbf{S}$
- B p
- C o
- D f

(Total for Question 6 = 1 mark)

- Ammonium iron(II) sulfate,  $(NH_4)_2Fe(SO_4)_2 \cdot 6H_2O$ , is a double salt that is used a start source of iron(II) ions.

  Formula mass of the double salt?  $S = 32.1 \quad Fe = 55.8$ (1)

- X 284.0
- **C** 392.0
- X 447.8
- (b) Ammonium sulfate is used in the preparation of the double salt.

What types of bond are present in ammonium sulfate?

(1)

- **A** ionic only
- X covalent and ionic only
- X **C** dative covalent and ionic only
- X **D** ionic, covalent and dative covalent
- (c) What is the **total** number of ions present in 0.1 mol of the double salt?

[Avogadro constant (L) =  $6.02 \times 10^{23} \text{ mol}^{-1}$ ]

(1)

- $1.80 \times 10^{23}$ X
- X **B**  $2.41 \times 10^{23}$
- **C**  $3.01 \times 10^{23}$ X
- X **D**  $6.62 \times 10^{23}$

(Total for Question 7 = 3 marks)

In which series are the elements in order of **increasing** melting temperature?  $D_{ij}$   $D_{$ 

X

X

X

X

(Total for Question 8 = 1 mark)

Which row gives the correct polarities of the S—F bond and the SF<sub>6</sub> molecule?

	Polarity of S—F bond	Polarity of SF <sub>6</sub> molecule
A	polar	polar
В	polar	non-polar
C	non-polar	polar
D	non-polar	non-polar

(Total for Question 9 = 1 mark)

- 10 Methane reacts with excess chlorine in UV light.
  - (a) Which process occurs in the initiation step?
    - A Cl
       Cl
    - B cl
       Cl

    - D Cl Cl
  - (b) Which of these molecules could **not** be formed in a termination step?

(1)

https://britishstatentro(1)

- $\square$  A  $C_2H_6$
- B CH<sub>3</sub>Cl
- C CH<sub>2</sub>Cl<sub>2</sub>
- D HCl

(Total for Question 10 = 2 marks)

- **11** Geometric isomerism is shown by 2-chlorobut-2-ene.
  - (a) What is the skeletal formula of *E*-2-chlorobut-2-ene?



- D
- (b) What is the **total** number of sigma bonds in Z-2-chlorobut-2-ene?

(1)

https://britishstudentroo(1)

- A 3
- B 4
- **C** 10

(Total for Question 11 = 2 marks)

**TOTAL FOR SECTION A = 20 MARKS** 

10

- 12 This question is about the chlorides of beryllium and calcium.
- SECTION B

  Answer ALL the questions. Write your answers in the spaces provided and the chlorides of beryllium and calcium.

  The of the atoms of beryllium and calcium

  (2) (a) Complete the electronic configurations of the atoms of beryllium and calcium using the s, p, d notation.



Be 1s<sup>2</sup>

Ca 1s<sup>2</sup>

(b) In the gaseous state, beryllium chloride is molecular.

Draw a dot-and-cross diagram to show the bonding in a molecule of beryllium chloride, BeCl<sub>2</sub>.

(2)

(c) In the solid state, beryllium chloride forms a polymeric structure.

The diagram shows part of this structure.

The diagram uses lines and arrows to represent the two different types of covalent bond.

Describe how each type of bond is formed.

(2)



رام) The Cl—Be—Cl bond angle is different in the two forms of beryllium chlorid	e.
(d) The Cl—Be—Cl bond angle is different in the two forms of beryllium chloridge.  Predict the two bond angles, justifying your answers by referring to electron-pair repulsion theory.	Astudentroomby Asia web
	. \$0 <sub>0</sub>
(e) Anhydrous calcium chloride is a crystalline, ionic solid which melts at 772°C.	
Draw a dot-and-cross diagram for calcium chloride. Show the outer electrons only.	
Show the outer elections only.	(2)



(f) Explain why gaseous beryllium chloride types of bonding.	and solid calcium chloride have different  (3)
	**de <sub>nfroon</sub> (3)
	ot the state of th
	(Total for Question 12 = 15 marks)



- 13 This question is about silicon and carbon.
  - (a) Silicon is a semiconductor.
    - (i) Data obtained using the mass spectrum of silicon are shown.

nicondu	con and carbon. uctor. ng the mass spectrum	of silicon are shown.	OS:/Britishstudentioon/b430a, Web. 400/
Iso	tope mass number	Relative abundance	The B. App.
	28	92.17	
	29	4.71	
	30	3.12	

Calculate the relative atomic mass of silicon to **two** decimal places.

(2)


(ii) Suggest a reason why there is a small peak in the mass spectrum of silicon at m/z = 14

(1)

(iii) Comp isotop	lete the table to show the of silicon.  Isotope  28Si	the number of protons	ماری and neutrons in each	Astuden,
	Isotope	Number of protons	Number of neutrons	Voon baso
	<sup>28</sup> Si			, Hell start
	<sup>29</sup> Si			
	<sup>30</sup> Si			

(1)

(b) Silicon dioxide, SiO<sub>2</sub>, is the main constituent of sand and has a giant lattice structure similar to that of diamond.



Crystalline silicon dioxide is used on the surface of semiconductor devices to provide a heat-resistant, electrically insulating layer.

Explain how the structure and bonding of silicon dioxide make it useful for this application.

(3)

Calcium silicate is formed in the removal of silicon dioxide impurited extraction of iron from its ores. A sample of calcium silicate composed of calcium, silicon and oxygen was found to contain 12.0 g of calcium, 8.43 g of silicon and oxygen.

14.47 g of oxygen. (c) Calcium silicate is formed in the removal of silicon dioxide impurities in the extraction of iron from its ores. A sample of calcium silicate composed of calcium,

(d) Carbon dioxide is a gas at room temperature. A fizzy drink is canned at 5.0 °C and  $1.3 \times 10^5$  Pa and contains approximately 3 g of carbon dioxide.

Calculate the volume, in cm<sup>3</sup>, occupied by 3.00 g of carbon dioxide gas at 5.0 °C and  $1.3 \times 10^5$  Pa.

$$[pV = nRT \quad R = 8.31 \,\mathrm{J}\,\mathrm{mol}^{-1}\,\mathrm{K}^{-1}]$$

(4)

(Total for Question 13 = 14 marks)



**14** Aluminium is an abundant metal with many uses.


(a) The first four ionisation energies of aluminium are shown.

undant metal with many use	es. um are shown.	Thips://Britishshidenfroomba30a.web.app
Ionisation number	Energy/kJ mol <sup>-1</sup>	Th. BARDA, L
1	578	*eb.
2	1820	
3	2750	
4	11600	

Explain how this information shows that aluminium is in Group 3 of the Periodic Table.

r	9	١.
	_//	
l.	4	

				j	
				7	
				_	
S	k	d	k	í	
S	Ŀ	2	k	7	
>			le N	2	
>			i	ĺ	
3					
3					
3			à		
>			à	5	
			à	5	
			à	5	
			à		
			à	5	
			à	5	
			à	5	
			à	5	
			à	5	
			à	5	
			à	5	
			à	5	
			à	5	
			à	5	
			à	5	
			à	5	
			à	5	
			à	5	
			à	5	
			à	5	
			à	5	



(i) Explain the general increase in the first ionisation energy from sodium to argon.

(2)

(ii) Explain why the first ionisation energy of aluminium is less than the first ionisation energy of magnesium.


(2)



(c) (i) Describe the bonding in aluminium metal.	(2)
(c) (i) Describe the bonding in aluminium metal.	Oom basila
(ii) Give two possible reasons why aluminium is used for overhead power cables.	(2)
(d) New uses for waste aluminium cans are being investigated. One possible use is to make nanoparticle alloys to produce hydrogen for fuel.	
(i) Aluminium nanoparticles react with water to produce aluminium oxide and hydrogen.	
Complete the following equation. State symbols are not required.	(1)
(ii) Give <b>two</b> possible reasons for producing hydrogen from aluminium rather	
than from fossil fuels.	(2)
(Total for Question 14 = 13 ma	rks)



<b>15</b> .	Alkanes	and	alkenes	are	obtained	from	crude	oil.
-------------	---------	-----	---------	-----	----------	------	-------	------



(b) (i) Octadecane can be cracked to produce butene and **one** other product. Complete the equation. State symbols are not required.

$$C_{18}H_{38} \rightarrow 2C_4H_8 +$$

(1)


(ii) One of the products of this cracking reaction is but-1-ene.

Give the **skeletal** formulae for the other three **alkene** isomers of  $C_4H_8$ 

(2)




(iii) Some reactions of but-1-ene are shown.



Give the name and **structural** formula of each of the products.

(4)

Product	Name	Structural formula
1		
2		
3		
4		



(d) State **one** advantage and **one** disadvantage of using incineration for the disposal of polymers, other than the effect on climate.

(2)

(e) (i) Butane is used as a fuel.

The equation for the complete combustion of butane is shown.

$$2C_4H_{10}(g) \ + \ 13O_2(g) \ \to \ 8CO_2(g) \ + \ 10H_2O(l)$$

35.0 cm<sup>3</sup> butane is completely burned in 300 cm<sup>3</sup> oxygen.

Calculate the final total volume of gas in cm<sup>3</sup>.

All volumes are measured at the same temperature and pressure.

(3)

				h <sub>the</sub>	
(ii)	Explain the main haza an enclosed space.	ard when using bu	utane as a fuel in a po	ortable hea <b>te</b> r in	
				Ode,	<b>3</b>
			(Total for Question 15 = 18 marks)		

TOTAL FOR SECTION B = 60 MARKS TOTAL FOR PAPER = 80 MARKS



### **Kr** krypton 36 **He** helium argon 18 131.3 **Xe** xenon [222] **Rn** radon 86 20.2 **Ne** ue 39.9 83.8 0 (8) 4.0 (18) Ar Elements with atomic numbers 112-116 have been reported 35.5 Cl chlorine fluorine 9 **Br** bromine 35 126.9 iodine 53 astatine 79.9 [210] / Selenium 34 127.6 **Te**tellurium 52 polonium oxygen 8 32.1 **S** sulfur [209] 79.0 16.0 but not fully authenticated 9 N | nitrogen | 7 As arsenic 33 121.8 **Sb** antimony hosphorus 209.0 Bi bismuth 31.0 **P** 74.9 വ germanium Carbon 28.1 **Si** silicon 14 72.6 **Sn** tin tin 50 207.2 g **Pb** lead 82 32 4 **Ga** gallium aluminium $\lim_{\text{indium}}$ thallium 114.8 27.0 10.8 **B** 204.4 69.7 49 m 200.6 **Hg** mercury 80 **Cd** cadmium 48 The Periodic Table of Elements **5.4 Zn** zinc 30 **Ag** silver 47 197.0 **Au** gold 79 **Cu** copper 29 [272] 195.1 Pt platinum 78 palladium darmstadtium 106.4 **Pd** [271] **Ds 58.7 Ni**nickel 46 Ir iridium rhodium 102.9 **Rh** 58.9 **Co** cobalt 27 192.2 [268] [98] 101.1 Tc Ru technetium ruthenium 1.0 **H** hydrogen 1 **Os** osmium [277] **HS** hassium 190.2 55.8 **Fe** iron 26 4 186.2 **Re** rhenium 75 [264] **Bh** bohrium 0 43 [266] **Sg** seaborgium molybdenum W tungsten chromium 183.8 95.9 Wo 42 atomic (proton) number relative atomic mass atomic symbol vanadium 23 **Ta** tantalum [262] **Db** dubnium niobium 180.9 92.9 **Nb** Key (2) rutherfordium zirconium titanium hafnium 178.5 [261] **Rf** 91.2 **Zr** 9 4 **La\*** lanthanum 45.0 Sc scandium 21 yttrium 39 actinium 138.9 88.9 [227] **Ac\*** $\widehat{\mathbb{C}}$ 22 Mg magnesium Calcium 20 strontium beryllium **Ba** barium 137.3 [226] **Ra** radium 87.6 **Sr** 24.3 40.1 9.0 38 K potassium 19 85.5 **Rb** rubidium **CS** caesium **Li** lithium rancium **Na** Sodium [223] **Fr** 132.9 23.0 39.1 22 6.9 37



<sup>\*</sup> Actinide series

hittps:	73 175 175 19 19 19 19 19 19 19 19 19 19 19 19 19	
	"den	
	175 <b>Lu</b> lutetium 71 os	[257] .a.o. <b>Lr</b> lawrencium 103
nticated	173 <b>Yb</b> ytterbium 70	No solution labelium la
but not fully authenticated	169 Tm thulium 69	[256] <b>Md</b> mendelevium 101
but not 1	167 <b>Er</b> erbium 68	[253] <b>Fm</b> fermium 100
	165 <b>Ho</b> holmium 67	[254] <b>Es</b> einsteinium 99
	163 <b>Dy</b> dysprosium 66	[251] Cf californium 98
roentgenium 111	159 <b>Tb</b> terbium 65	[245] <b>BK</b> berkelium 97
darmstadtium 110	157 <b>Gd</b> gadolinium 64	[247] <b>Cm</b> curium 96
meitnerium damstadtium roentgenium 109 110 111	152 <b>Eu</b> europium 63	[243]
hassium 108	150 Sm samarium 6	Pu Pu plutonium 94
bohrium 107	Pm promethium 61	[237] <b>Np</b> neptunium 93
seaborgium 106	144 <b>Nd</b> neodymium 60	238 U uranium 92
ntherfordium dubnium seaborgium	141 Pr praseodymium 59	[231] <b>Pa</b> protactinium 91
rutherfordium 104	140 <b>Ce</b> cerium 58	232 <b>Th</b> thorium 90
۶		

