

Please check the examination details below before entering your candidate information

Candidate surname

Other names

Centre Number

Candidate Number

--	--	--	--	--

--	--	--	--

## Pearson Edexcel International Advanced Level

Time 1 hour 30 minutes

Paper  
reference

**WME01/01**

### Mathematics

### International Advanced Subsidiary/Advanced Level Mechanics M1

**You must have:**

Mathematical Formulae and Statistical Tables (Yellow), calculator

Total Marks

**Candidates may use any calculator permitted by Pearson regulations. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.**

#### Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided  
– *there may be more space than you need.*
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Whenever a numerical value of  $g$  is required, take  $g = 9.8 \text{ m s}^{-2}$ , and give your answer to either 2 significant figures or 3 significant figures.

#### Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 8 questions in this question paper. The total mark for this paper is 75.
- The marks for **each** question are shown in brackets  
– *use this as a guide as to how much time to spend on each question.*

#### Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.
- If you change your mind about an answer, cross it out and put your new answer and any working underneath.

Turn over ►

P66649A


©2021 Pearson Education Ltd.

E:1/1/1/



  
Pearson

1.



**Figure 1**

A non-uniform rod  $AB$  has length  $9\text{ m}$  and mass  $M\text{ kg}$ .

The rod rests in equilibrium in a horizontal position on two supports, one at  $C$  where  $AC = 2.5\text{ m}$  and the other at  $D$  where  $DB = 2\text{ m}$ , as shown in Figure 1.

The magnitude of the force acting on the rod at  $D$  is twice the magnitude of the force acting on the rod at  $C$ .

The centre of mass of the rod is  $d$  metres from  $A$ .

Find the value of  $d$ .

**(6)**

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA





Question 1 continued

Lined area for writing the answer to Question 1.

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA





2. A particle  $P$  of mass  $2m$  is moving on a rough horizontal plane when it collides directly with a particle  $Q$  of mass  $4m$  which is at rest on the plane. The speed of  $P$  immediately before the collision is  $3u$ . The speed of  $Q$  immediately after the collision is  $2u$ .

(a) Find, in terms of  $u$ , the speed of  $P$  immediately after the collision. (3)

(b) State clearly the direction of motion of  $P$  immediately after the collision. (1)

Following the collision,  $Q$  comes to rest after travelling a distance  $\frac{6u^2}{g}$  along the plane.

The coefficient of friction between  $Q$  and the plane is  $\mu$ .

(c) Find the value of  $\mu$ . (6)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



### Question 2 continued

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA







Question 2 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Lined writing area for question 2.

(Total 10 marks)

Q2

Marking boxes for Q2.



3. A car is moving at a constant speed of  $25 \text{ m s}^{-1}$  along a straight horizontal road.

The car is modelled as a particle.

At time  $t = 0$ , the car is at the point  $A$  and the driver sees a road sign  $48 \text{ m}$  ahead.

Let  $t$  seconds be the time that elapses after the car passes  $A$ .

In a **first** model, the car is assumed to decelerate uniformly at  $6 \text{ m s}^{-2}$  from  $A$  until the car reaches the road sign.

(a) Use this first model to find the speed of the car as it reaches the sign. (2)

The road sign indicates that the speed limit immediately after the sign is  $13 \text{ m s}^{-1}$ .

In a **second** model, the car is assumed to decelerate uniformly at  $6 \text{ m s}^{-2}$  from  $A$  until it reaches a speed of  $13 \text{ m s}^{-1}$ . The car then maintains this speed until it reaches the road sign.

(b) Use this second model to find the value of  $t$  at which the car reaches the sign. (4)

In a **third** model, the car is assumed to move with constant speed  $25 \text{ m s}^{-1}$  from  $A$  until time  $t = 0.2$ , the car then decelerates uniformly at  $6 \text{ m s}^{-2}$  until it reaches a speed of  $13 \text{ m s}^{-1}$ . The car then maintains this speed until it reaches the road sign.

(c) Use this third model to find the value of  $t$  at which the car reaches the sign. (4)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

DO NOT WRITE IN THIS AREA







**Question 3 continued**

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

**Q3**

**(Total 10 marks)**



P 6 6 6 4 9 A 0 1 3 3 2

4. The position vector,  $\mathbf{r}$  metres, of a particle  $P$  at time  $t$  seconds, relative to a fixed origin  $O$ , is given by

$$\mathbf{r} = (t - 3)\mathbf{i} + (1 - 2t)\mathbf{j}$$

- (a) Find, to the nearest degree, the size of the angle between  $\mathbf{r}$  and the vector  $\mathbf{j}$ , when  $t = 2$  (3)
- (b) Find the values of  $t$  for which the distance of  $P$  from  $O$  is 2.5 m. (5)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA











5.




Figure 2

A small bead of mass 0.2 kg is attached to the end  $P$  of a light rod  $PQ$ . The bead is threaded onto a fixed vertical rough wire.

The bead is held in equilibrium with the rod  $PQ$  inclined to the wire at an angle  $\alpha$ , where  $\tan \alpha = \frac{4}{3}$ , as shown in Figure 2.

The thrust in the rod is  $T$  newtons.

The bead is modelled as a particle.

- (a) Find the magnitude and direction of the friction force acting on the bead when  $T = 2.5$  (3)

The coefficient of friction between the bead and the wire is  $\mu$ .

Given that the greatest possible value of  $T$  is 6.125

- (b) find the value of  $\mu$ . (7)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA





**Question 5 continued**

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Lined writing area for the answer to Question 5.









7.




Figure 4

One end of a light inextensible string is attached to a particle  $A$  of mass  $2m$ . The other end of the string is attached to a particle  $B$  of mass  $3m$ . The string passes over a small, smooth, light pulley  $P$  which is fixed at the top of a rough inclined plane. The plane is inclined to the horizontal at an angle  $\alpha$ , where  $\tan \alpha = \frac{3}{4}$

Particle  $A$  is held at rest on the plane with the string taut and  $B$  hanging freely below  $P$ , as shown in Figure 4. The section of the string  $AP$  is parallel to a line of greatest slope of the plane.

The coefficient of friction between  $A$  and the plane is  $\frac{1}{2}$

Particle  $A$  is released and begins to move up the plane.

For the motion before  $A$  reaches the pulley,

- (a) (i) write down an equation of motion for  $A$ ,
  - (ii) write down an equation of motion for  $B$ ,
- (4)**
- (b) find, in terms of  $g$ , the acceleration of  $A$ ,
- (5)**
- (c) find the magnitude of the force exerted on the pulley by the string.
- (4)**
- (d) State how you have used the information that  $P$  is a smooth pulley.
- (1)**

---

---

---

---

---

---

---

---

---

---

DO NOT WRITE IN THIS AREA





Question 7 continued

Lined area for writing the answer to Question 7.

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Leave  
blank

### Question 7 continued

Handwritten response area consisting of 30 horizontal lines.

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA













