| Please check the examination det | ails bel | ow before ente | ering your candidate information |
|---|----------|--------------------|----------------------------------|
| Candidate surname | | | Other names Ohiri |
| Pearson Edexcel International GCSE (9–1) | Cen | tre Number | Candidate Number |
| Time 2 hours | | Paper reference | ADH1/1D ASDO/1D |
| Physics | | | |
| Science (Double Award PAPER: 1P | d) 49 | SD0 | |
| | | | |
| You must have: Calculator, ruler, protractor | | | Total Marks |

Instructions

- Use **black** ink or ball-point pen.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Answer the questions in the spaces provided
 - there may be more space than you need.
- Show all the steps in any calculations and state the units.
- Some questions must be answered with a cross in a box ⋈. If you change your mind about an answer, put a line through the box ⋈ and then mark your new answer with a cross ⋈.

Information

- The total mark for this paper is 110.
- The marks for **each** question are shown in brackets
 - use this as a guide as to how much time to spend on each question.

Advice

- Read each question carefully before you start to answer it.
- Write your answers neatly and in good English.
- Try to answer every question.
- Check your answers if you have time at the end.
- Good luck with your examination.

Turn over ▶







http://britishstatentroom.worthress.com/

FORMULAE

You may find the following formulae useful.

energy transferred = $current \times voltage \times time$

$$E = I \times V \times t$$

frequency =
$$\frac{1}{\text{time period}}$$

$$f = \frac{1}{T}$$

$$power = \frac{work done}{time taken}$$

$$P = \frac{W}{t}$$

$$power = \frac{energy transferred}{time taken}$$

$$P = \frac{W}{t}$$

orbital speed =
$$\frac{2\pi \times \text{orbital radius}}{\text{time period}}$$

$$v = \frac{2 \times \pi \times r}{T}$$

(final speed)² = (initial speed)² + $(2 \times acceleration \times distance moved)$

$$v^2 = u^2 + (2 \times a \times s)$$

$$pressure \times volume = constant$$

$$p_1 \times V_1 = p_2 \times V_2$$

$$\frac{\text{pressure}}{\text{temperature}} = \text{constant}$$

$$\frac{p_1}{T_1} = \frac{p_2}{T_2}$$

Where necessary, assume the acceleration of free fall, $g = 10 \text{ m/s}^2$.

http://britishstudentroom.wordpress.com/

BLANK PAGE

Answer ALL questions.

- hito: Britishstatentroom, wordpress.com (a) (i) Which of these coloured stars has the highest surface temperature?
 - orange X
 - red
 - white X
 - **D** yellow
 - (ii) Which of these is a stage in the life cycle of the Sun?

(1)

- black hole
- neutron star
- supernova
- white dwarf
- (iii) Which of these is the stage nearest the end of the life cycle of a star with a mass much greater than the Sun?

(1)

- main sequence
- protostar
- C supernova
- white dwarf

(b) The diagram shows the orbit of a comet around a star.

Draw a labelled arrow to show the force acting on the comet due to the star.

not to scale

comet

(Total for Question 1 = 5 marks)



- A student uses iron filings to investigate the magnetic field pattern around a bar magnet.


 (a) Name the apparatus that the student could use to find the direction of the field lines.

 (b) The could be apparatus that the student could use to find the direction of the field lines.

The student makes three mistakes.

Draw a circle around each mistake on the diagram.

(3)



| | Ath. |
|---|--|
| (c) The magnet is made of a magnetically hard material. | v. British |
| Describe what is meant by the term magnetically ha | h _{th./britishstudentro(2)} ard. |
| | To the state of th |
| | |
| | (Total for Question 2 = 6 marks) |
| | |


(1)

The diagram shows a model of a chain reaction in a nuclear fission reactor.

A ball falls through the hole and hits a mousetrap.

This mousetrap then releases another ball.

Each released ball hits a different mousetrap.




- (a) Each mousetrap represents a uranium-235 nucleus.
 - (i) Name the particle represented by the balls.

(ii) Describe what is meant by the term **nuclear fission**. (2)

| (111) | Describe what is | meant by the | term chain reacti | on. | Britishs _k | (2) |
|-----------------------------|---|----------------------------------|-----------------------------------|-------------------------------|-----------------------|-----------|
| | | | | | htp://britishstude | Atroop |
| | | | | | | V. Wordpr |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| b) The | walls of a real n | uclear reactor a | re not made of pl | astic. | | |
| Give | e a suitable mate | erial for the wal | ls of a real nuclear | r reactor. | | (1) |
| | | | | | | |
| | | | | | | |
| c) Use | words from the | box to comple | te the sentences a | about the mod | erator and | |
| | words from the trol rods in a nuc | • | te the sentences a | about the mode | erator and | (4) |
| | | • | te the sentences a | about the mod | erator and | (4) |
| | | • | te the sentences a | about the mode | erator and slow | (4) |
| | trol rods in a nuc | clear reactor. | | | | (4) |
| con | absorb boron | ionise granite | neutralise graphite | reflect paper | slow uranium | (4) |
| modera | absorb boron ator in a nuclear | ionise granite reactor can be | neutralise graphite made of | reflect paper | slow uranium | (4) |
| modera | absorb boron ator in a nuclear | ionise granite reactor can be | neutralise graphite made of | reflect paper | slow uranium | |
| modera | absorb boron ator in a nuclear | ionise granite reactor can be | neutralise graphite made of | reflect paper | slow uranium | |
| modera modera control | absorb boron ator in a nuclear ator is designed | ionise granite reactor can be to | neutralise graphite made of | reflect paper neutrons. | slow uranium | |





Time in seconds

(a) (i) Calculate the acceleration of the car between 1.8 and 8.0 seconds.

(3)

(ii) Calculate the braking distance of the car.


(3)


| (iii) Explain the effect, if any, of increased driver tiredness on the thinking | distance |
|---|-------------------|
| and on the braking distance of the car. | *Sh _{Sh} |


braking distance......


(b) Which of these represents the distance-time graph for the car?

(1)










(Total for Question 4 = 11 marks)

http://britishstudentroom.worthress.com/

BLANK PAGE



(a) Draw meters on the diagram to measure the voltage of the LED and the current in the LED. (3)

(b) The table shows the results of the investigation.

| Voltage in V | Current in mA |
|--------------|---------------|
| 0.00 | 0.0 |
| 0.30 | 0.5 |
| 0.35 | 2.5 |
| 0.40 | 1.5 |
| 0.45 | 2.0 |
| 0.50 | 4.5 |
| 0.55 | 9.0 |
| 0.60 | 15.0 |

Alto: Abilish studentroom, worth ress. com

(i) Plot the results on the grid.

(3)


(ii) One of the results is anomalous.

On the graph, draw a circle around the anomalous result.

(1)

(iii) Draw a curve of best fit.

(1)




14

| (iv) Give a reason why a line graph is the best way of | ~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~ |
|--|--|
| | |
| (v) State the formula linking voltage, current and re | sistance. |
| (vi) Any current larger than 15 mA will norman enthy | damage the LED |
| (vi) Any current larger than 15 mA will permanently The resistor in the circuit has a resistance of 270 | _ |
| | |
| Use the results from the investigation to determ the power supply without damaging the LED. | |
| | (4 |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| ma | ximum voltage = |
| | Total for Question 5 = 14 marks |
| | iotal for Question 5 = 14 marks |

or and a lid. The temperature on sensor The diagram shows a container of water. The container has a heater in the bottom corner, a temperature sensor and a lid. lid heater (a) Explain how a convection current is formed in the container. You may add to the diagram or draw your own diagram to support your answer. (4)

(b) The graph shows how the temperature sensor reading changes with time. The street of the change in temperature when the container is full of water and when the container is empty.



is full of water.

Key
---- container empty
---- container full


(i) Describe the relationship between temperature and time when the container

(2)

(ii) Explain the differences between the two curves on the graph.

(3)

(Total for Question 6 = 9 marks)



Alpha particles produce the shorter, thicker tracks.

Beta particles produce the longer, thinner tracks.

Explain why alpha particles produce the shorter, thicker tracks.


(2)

- (b) Uranium-235 is an isotope of uranium that can decay by emitting an alpha particle.
 - (i) Describe what is meant by the term **isotope**.

(2)



(ii) Complete the equation for the decay of uranium-235



(iii) Uranium-235 decays to thorium-231 with a half-life of 700 million years.

When a rock was formed, it contained 6400 million uranium-235 nuclei and no thorium-231 nuclei.

Show that after 2100 million years there are seven times more thorium nuclei than uranium nuclei in the rock.

(5)


(Total for Question 7 = 11 marks)

(a) Describe how two magnets can be arranged to produce a uniform magnetic field.

You may draw a diagram to support your answer.

**The production of the product of the p

(b) The diagram shows a wire in a uniform magnetic field.



(i) The wire is moved up and down repeatedly in the magnetic field. Explain the effect that this movement has on the ammeter reading.

(3)



| (ii) Explain the effects on the ammeter reading amplitude but at a higher frequency. | of moving the wire with the same |
|--|----------------------------------|
| | Wycess Com |
| | |
| | |
| | |
| | |
| | (Total for Question 8 = 9 marks) |

9 The photograph shows a whale jumping out of the surface of the sea.



Alto: Abilish studentroom, worth ress. com

(Source: © Alexander Baumann/Shutterstock)

(a) At the top of the jump, the whale's velocity is 0 m/s.

The whale falls 2.2 m from the top of the jump to the surface of the sea.

Calculate the velocity of the whale when it hits the surface of the sea.

(4)

velocity = m/s

- (b) A resultant force causes the whale to slow down when it hits the surface of the sea.

 "` Praw an arrow to show this resultant force.

 "The sea of the sea.

 "The sea of the sea of the sea of the sea.

 "The sea of the sea of the



(Source: © Eugenia Petrovskaya/Shutterstock)

- (ii) The resultant force acting on the whale is 18000 N.
 - The mass of the whale is 4100 kg.
 - Calculate the acceleration of the whale.

(3)


acceleration = m/s²

(Total for Question 9 = 8 marks)



10 The diagram shows a device called a hydraulic lift.

The hydraulic lift consists of a tube of oil with a piston at each end.



(a) Calculate the pressure difference between the bottom of piston X and the bottom of the oil.

[density of oil = 820 kg/m^3]

pressure difference =Pa

- (b) A 24 kg mass is placed on piston X.
 - (i) Calculate the weight of the 24 kg mass.

Give the unit.

(3)

(3)

weight = unit

| (| (ii) | Calculate | the extra | pressure o | n the oi | il due to | the mass |
|---|--------------|-----------|-----------|------------|----------|-----------|----------------|
| ١ | (II <i>)</i> | Calculate | uie exua | Diessule 0 | n the o | ii uue to | ı tile illass. |

| Ath. Brit. | |
|------------------------|-------------|
| ish _{shude} , | (2) |
| ** | Itroopp. W. |
| | oridoriese |
| | , com |

extra pressure =Pa

(iii) The oil transfers the same extra pressure to piston Y.

Calculate the force acting upwards on piston Y due to the extra pressure.

(3)

force = N

(c) Piston Y starts at rest, rises slowly and then comes to rest.

State how the following energy stores have changed from before the motion to after the motion is complete.

(4)

| Energy store | Change |
|--|--------|
| gravitational potential energy of piston X | |
| gravitational potential energy of piston Y | |
| chemical energy of piston Y | |
| kinetic energy of piston Y | |

(Total for Question 10 = 15 marks)




Diagram 1

- (a) When very bright light shines on this transparent material, a small amount of light is reflected at point X.
 - (i) On diagram 2, draw the reflected ray from point X.




Diagram 2

(ii) Measure the angle of **refraction** at point X on diagram 2.

(1)

(2)

angle =degrees

(iv) Show that the refractive index of the transparent material is about 1.7

(2)

(v) State the formula linking refractive index and critical angle.

(1)


(vi) Calculate the critical angle for the transparent material.

(2)

critical angle =degrees

QUESTION 11 CONTINUES ON NEXT PAGE





TOTAL FOR PAPER = 110 MARKS